



## Contribution to Special Issue: 'Towards a Broader Perspective on Ocean Acidification Research' Original Article

# Ocean warming and elevated carbon dioxide: multiple stressor impacts on juvenile mussels from southern Chile

Jorge M. Navarro<sup>1,2\*</sup>, Cristian Duarte<sup>3,4</sup>, Patricio H. Manríquez<sup>5</sup>, Marco A. Lardies<sup>6</sup>, Rodrigo Torres<sup>7</sup>, Karin Acuña<sup>1</sup>, Cristian A. Vargas<sup>8</sup>, and Nelson A. Lagos<sup>9</sup>

<sup>1</sup>Instituto de Ciencias Marinas y Limnológicas, Laboratorio Costero de Calfuco, Facultad de Ciencias, Universidad Austral de Chile, Valdivia, Chile

<sup>2</sup>Centro Fonmap de Investigación de Ecosistemas Marinos de Altas Latitudes (IDEAL), Valdivia, Chile

<sup>3</sup>Departamento de Ecología y Biodiversidad, Facultad de Ecología y Recursos Naturales, Universidad Andres Bello, Santiago, Chile

<sup>4</sup>Center for the Study of Multiple-drivers on Marine Socio-Ecological System (MUSELS), Universidad de Concepción, Concepción, Chile

<sup>5</sup>Laboratorio de Ecología y Conducta de la Ontogenia Temprana (LECOT), Centro de Estudios Avanzados en Zonas Áridas (CEAZA), Av. Ossandón 877, Coquimbo, Chile

<sup>6</sup>Facultad de Artes Liberales and Facultad de Ingeniería y Ciencias, Universidad Adolfo Ibáñez, Santiago, Chile

<sup>7</sup>Centro de Investigación en Ecosistemas de la Patagonia (CIEP), Coyhaique, Chile

<sup>8</sup>Laboratorio de Funcionamiento de Ecosistemas Acuáticos (LAFE), Departamento de Sistemas Acuáticos, Facultad de Ciencias Ambientales, Universidad de Concepción, Concepción, Chile

<sup>9</sup>Centro de Investigación e Innovación para el Cambio Climático, Universidad Santo Tomás, Santiago, Chile

\*Corresponding author: tel: + 56 63 2221556; fax: + 56 63 2221455; e-mail: [jnavarro@uach.cl](mailto:jnavarro@uach.cl)

Navarro, J. M., Duarte, C., Manríquez, P. H., Lardies, M. A., Torres, R., Acuña, K., Vargas, C. A., and Lagos, N. A. Ocean warming and elevated carbon dioxide: multiple stressor impacts on juvenile mussels from southern Chile. – ICES Journal of Marine Science, 73: 764–771.

Received 17 June 2015; revised 24 November 2015; accepted 25 November 2015; advance access publication 4 January 2016.

The combined effect of increased ocean warming and elevated carbon dioxide in seawater is expected to have significant physiological and ecological consequences at many organizational levels of the marine ecosystem. In the present study, juvenile mussels *Mytilus chilensis* were reared for 80 d in a factorial combination of two temperatures (12 and 16°C) and three  $p\text{CO}_2$  levels (380, 700, and 1000  $\mu\text{atm}$ ). We investigated the combined effects of increasing seawater temperature and  $p\text{CO}_2$  on the physiological performance (i.e. feeding, metabolism, and growth). Lower clearance rate (CR) occurred at the highest  $p\text{CO}_2$  concentration (1000  $\mu\text{atm}$ ) compared with the control (380  $\mu\text{atm}$ ) and with the intermediate concentration of  $p\text{CO}_2$  (700  $\mu\text{atm}$ ). Conversely, CR was significantly higher at 16°C than at 12°C. Significant lower values of oxygen uptake were observed in mussels exposed to 1000  $\mu\text{atm}$   $p\text{CO}_2$  level compared with those exposed to 380  $\mu\text{atm}$   $p\text{CO}_2$ . Scope for growth (SFG) was significantly lower at the highest  $p\text{CO}_2$  concentration compared with the control. Mussels exposed to 700  $\mu\text{atm}$   $p\text{CO}_2$  did not show significantly different SFG from the other two  $p\text{CO}_2$  treatments. SFG was significantly higher at 16°C than at 12°C. This might be explained because the experimental mussels were exposed to temperatures experienced in their natural environment, which are within the range of thermal tolerance of the species. Our results suggest that the temperature rise within the natural range experienced by *M. chilensis* generates a positive effect on the processes related with energy gain (i.e. feeding and absorption) to be allocated to growth. In turn, the increase in the  $p\text{CO}_2$  level of 1000  $\mu\text{atm}$ , independent of temperature, adversely affects this species, with significantly reduced energy allocated to growth (SFG) compared with the control treatment.

**Keywords:** high  $\text{CO}_2$ , multiple stressors, mussels, ocean warming, scope for growth, thermal window.

## Introduction

Fossil fuel burning and land use change are the dominant causes of the increase in atmospheric  $\text{CO}_2$  concentration from pre-industrial

levels (ca. 280  $\mu\text{atm}$ ) reaching current values of ca. 400  $\mu\text{atm}$  (IPCC, 2013). Atmospheric  $\text{CO}_2$  concentration increased at an average rate of  $2.0 \pm 0.1 \mu\text{atm year}^{-1}$  during 2002–2011, being higher than any

previous decade since direct atmospheric concentration measurements began in 1958 (IPCC, 2013). According to the current tendency, it is expected that by the end of this century atmospheric CO<sub>2</sub> concentration could reach 700  $\mu\text{atm}$  and will exceed 1500  $\mu\text{atm}$  between the years 2100 and 2200 (Caldeira and Wickett, 2003; Pörtner et al., 2004; IPCC, 2013). Approximately 30% of atmospheric CO<sub>2</sub> dissolves in the ocean, resulting in reduction of pH and the availability of carbonate ions, which are essential to calcium carbonate deposition in calcifier organisms. According to the current trend of increasing CO<sub>2</sub> concentration, ocean acidification (OA) will decrease from 0.2 to 0.4 pH units by 2100, together with a reduction of 50% of carbonate ions (Caldeira and Wickett, 2003; Feely et al., 2004). OA is not an isolated stressor, yet acts together with other environmental shifts such as ocean warming (Meier, 2006; IPCC, 2013). Sea surface temperature has increased by  $\sim 0.6^\circ\text{C}$  during the last 50 years (Walther et al., 2002), and it is expected that it will continue to rise between 1 and  $4^\circ\text{C}$  by the end of the century (IPCC, 2013).

Growing evidence shows that many environmental stressors can act synergistically, affecting many physiological processes of marine organisms (Schiedek et al., 2007; Gooding et al., 2009). Increased CO<sub>2</sub> concentration in seawater has significant physiological and ecological consequences at many organizational levels of the marine ecosystem (Fabry et al., 2008). A wide range of sensitivities to projected rates of OA exist within and across groups of marine organisms (e.g. corals, oysters, mussels, crabs, and sea urchins), with a trend for greater sensitivity in early life stages (Vargas et al., 2013; Gobler and Talmage, 2014). However, key uncertainties remain in our understanding of the impacts on organisms, life histories, and ecosystems. Most results describe that marine organisms are affected by high  $p\text{CO}_2$  levels by disturbances in acid–base regulation, respiration, metabolism, growth rates, reproduction, and calcification (Pörtner, 2008; Widdicombe and Spicer, 2008; Navarro et al., 2013; Duarte et al., 2014; Gazeau et al., 2014).

Because many marine organisms live close to their thermal compensatory capacity (Somero, 2002), the increase in temperature associated with global climate change is expected to impact all physiological processes, survival, and many ecological interactions (Godbold and Solan, 2013). Biological response to rising temperature may vary within and among species, and even across ontogenetic stages (Harley et al., 2006). Seawater temperature is shown to have strong effects on the physiological performance (Pörtner, 2010; Hiebenthal et al., 2013) and reproduction (Navarro et al., 2000; Philippart et al., 2003) of benthic organisms. Recent studies show that the effects of increased CO<sub>2</sub> could be modified by a rise in temperature (Gooding et al., 2009; Findlay et al., 2010). The interaction of elevated seawater CO<sub>2</sub> and extreme temperatures is described to narrow the thermal tolerance window of an organism exposed to high CO<sub>2</sub> levels (Pörtner, 2008; Pörtner and Farrell, 2008). Metzger et al. (2007) found that at elevated seawater CO<sub>2</sub> concentrations, the upper thermal tolerance limits were reduced by several Celsius degrees in crustaceans. High temperature and elevated CO<sub>2</sub> lead to a significant decrease in shell hardness in the oyster *Crassostrea virginica* and the clam *Mercenaria mercenaria*, suggesting major changes in their biomineralization processes (Ivanina et al., 2013). However, early juveniles and adults may be vulnerable to skeletal dissolution, although warming may diminish the negative impact of acidification on calcification (Byrne, 2011). In a recent study, we found that the net rate of calcium deposition in juveniles of *Mytilus chilensis* was not significantly affected by temperature, but was negatively affected by high levels of CO<sub>2</sub> (Duarte et al., 2014).

In high-latitude coastal areas of South America, the edible mussel *M. chilensis* forms extensive subtidal and intertidal beds that play an important ecological role, affecting the community structure of the associated macrofauna (Duarte et al., 2006). *Mytilus chilensis* is an important species in Chile for aquaculture activities, with a landing of ca. 300 000 tons per year. These facts make this bivalve species a key group of marine invertebrates for studying the biological impacts of the interaction of elevated  $p\text{CO}_2$  levels and global warming on their physiology and growth. In a previous study, Navarro et al. (2013) described the effects of high levels of  $p\text{CO}_2$  for another population of *M. chilensis* of southern Chile, concluding that growth rate might be significantly reduced under medium-term exposure to elevated  $p\text{CO}_2$  levels.

In the present study, juvenile individuals of the mussel *M. chilensis* were reared in a factorial combination of two temperatures (12 and  $16^\circ\text{C}$ ) and three  $p\text{CO}_2$  levels (380, 700, and 1000  $\mu\text{atm}$ ). We analysed the combined effects of increasing seawater temperature and  $p\text{CO}_2$  on the physiological performance (i.e. feeding, metabolism, and growth) of *M. chilensis*. We hypothesized that the sensitivity of juvenile mussel *M. chilensis* exposed to elevated levels of  $p\text{CO}_2$  increases with rising temperature.

## Material and methods

Juvenile individuals of *M. chilensis* were collected from suspended cultures (permanently submerged) at Huelmo Bay, Puerto Montt ( $41^\circ 67'\text{S}$ ,  $73^\circ 03'\text{W}$ ), during the 2011 winter season (sea surface temperature =  $11.5^\circ\text{C}$ ; pH 8.09 at the day of collection) and transported to the laboratory under chilled conditions. Mussels of approximately equal length ( $24.2 \pm 0.4$  mm SE) and dry tissue weight ( $33.4 \pm 1.6$  mg SE) were selected. Before the experiments, mussels were acclimated for 2 weeks in aquaria with circulating seawater (temperature =  $12\text{--}13^\circ\text{C}$  and salinity 32–33 psu) and fed daily with the microalgae *Isochrysis galbana*.

## Experimental set-up

Following the acclimation period, experimental mussels were transferred to the aquaria and exposed to six different treatments at the combination of two temperatures (12 and  $16^\circ\text{C}$ ) and three  $p\text{CO}_2$  levels (380, 700, and 1000  $\mu\text{atm}$ ), for a period of 80 d. Each treatment was replicated four times, and each replicate contained five experimental animals, which were identified using bee tags. The experimental temperatures were within the range of thermal tolerance of the species;  $12^\circ\text{C}$  represents the average and  $16^\circ\text{C}$  the extreme highest value for the area although this was last recorded in a narrow temporal window during summer (Navarro and Jaramillo, 1994). Experimental  $p\text{CO}_2$  concentrations were selected to match current day and expected near-future levels according to the predicted values by the IPCC (2013), based on the extreme scenario (RCP8.5) of atmospheric CO<sub>2</sub>. Two thermo-regulated baths were prepared for each experimental temperature (12 and  $16^\circ\text{C}$ ) consisting each of 12 plastic 3.5 l tanks (four replicates for each  $p\text{CO}_2$  treatment). The different  $p\text{CO}_2$  concentrations were obtained using a mesocosm system, following Navarro et al. (2013) and Torres et al. (2013). Briefly, for 380  $\mu\text{atm}$ , pure atmospheric air was bubbled into experimental containers. For 700 and 1000  $\mu\text{atm}$ , dry air was blended with pure CO<sub>2</sub> to each CO<sub>2</sub> concentration using mass flow controllers (MFCs, [www.aalborg.com](http://www.aalborg.com)) for air and CO<sub>2</sub>. During the experiments, water pH and total alkalinity were monitored every 3 d in equilibration tanks. pH measurements were done in closed 25 ml cells thermostated at  $25^\circ\text{C}$  using a Metrohm 713 pH meter, calibrated with 8.089 Tris-buffer at  $25^\circ\text{C}$ . Temperature and salinity

were measured in the equilibration tanks using a small CTD (Ocean Seven 305 Plus CTD, [www.lidronaut.it](http://www.lidronaut.it)). Total alkalinity was measured using the method of Haraldsson *et al.* (1997). Resulting hydrographic data were used to calculate the rest of the carbonate system parameters and the saturation stage of Omega Aragonite and Calcite using CO<sub>2</sub>SYS software (Lewis and Wallace, 1998) set with Mehrbach solubility constants (Mehrbach *et al.*, 1973) refitted by Dickson and Millero (1987). All experimental individuals were fed with a daily amount of food (*I. galbana*) equal to 5% of the body weight. An open system, consisting of a stock suspension of the microalgae and a multichannel peristaltic pump (Masterflex 7524), was used to deliver the suspension directly to each experimental aquarium. Food concentration was maintained within a range similar to that found in the natural southern Chilean environment throughout the year (ca. 0.8–1.5 mg l<sup>-1</sup> dry weight). Seawater was gently changed every day, with the corresponding pCO<sub>2</sub> level from the seawater acidification system.

### Physiological parameters

Physiological measurements (below) were done on 120 individual mussels taken randomly from each experimental aquarium of the six treatments (two temperatures and three pCO<sub>2</sub> concentrations), starting on day 0 (initial exposure) and continuing every 20 d along a period of 80 d.

### Feeding experiments

Clearance rate (CR) of mussels exposed to the experimental temperature and CO<sub>2</sub> combinations (six treatments) was estimated in a static system homogenized by aeration using a food concentration of  $25 \times 10^6$  cells *I. galbana* l<sup>-1</sup>. One juvenile mussel was taken randomly from the replicates of each treatment and placed in an experimental chamber (0.4 l). Subsequently, the decrease in the number of particles was monitored every 30 min over a period of 3 h, using an Elzone 180XY particle counter, equipped with a tube of 120- $\mu$ m aperture. A control aquarium without mussels was used to discard the sedimentation of particles and the CR (l h<sup>-1</sup> mussel<sup>-1</sup>) was calculated according to Coughlan (1969). Organic ingestion rate was calculated as the product of the CR and the organic material contained in the diet.

After CR measurements were completed, absorption efficiency (AE) was estimated by determining the organic and inorganic content of the food and the faeces following the ratio method of Conover (1966). Samples of food and faeces were filtered through pre-washed, preweighed, 25-mm glassfibre filters. Filters were rinsed with isotonic ammonium formate, dried to a constant weight at 100°C, weighed, combusted at 450°C for 3 h, and weighed again to estimate the organic and inorganic fraction contained in the food and faeces. Absorption rate was calculated as the product of the organic ingestion rate and AE.

### Ammonia excretion and oxygen uptake

Ammonia excretion and oxygen uptake were determined immediately after the CR measurements were made. Individual mussels were placed in sealed glass beakers (0.14 l) containing filtered (0.45  $\mu$ m) seawater. One additional beaker containing filtered seawater but without mussels was used as a control. All beakers were filled with the corresponding experimental pCO<sub>2</sub> levels (380, 700, and 1000  $\mu$ atm) and maintained at 12 and 16°C by submerging them in two thermostatic water baths. After 2 h, water samples (5 ml) from each beaker were taken and analysed for ammonia–nitrogen by the phenol-hypochlorite colorimetric method of

Solórzano (1969). Values for excretion rate were expressed in microgram of NH<sub>4</sub>-N. The oxygen uptake by each mussel was estimated by the difference between the oxygen contained in the control and that contained in the experimental beaker over the period of 2 h and analysed by the micro-Winkler method, modified by Ohle (1953), which uses titration to determine dissolved oxygen in the water.

### Scope for growth

Scope for growth (SFG) was calculated after converting all the physiological rates to energy equivalents (J h<sup>-1</sup>): 1 ml O<sub>2</sub> = 19.9 J; 1  $\mu$ g NH<sub>4</sub>-N = 0.0249 J (Elliot and Davison, 1975), and 1 mg of organic material of *I. galbana* = 18.75 J (Whyte, 1987).

### Statistical analyses

To avoid pseudo-replication problems, the measured physiological processes were averaged for five mussels of each replicate. A two-way ANOVA was used to estimate the effects of temperature and pCO<sub>2</sub> concentrations on different physiological measurements. When the analyses did not show significant interactions, multiple comparisons were carried out using Tukey's *a posteriori* HSD test on each factor that showed significant differences (Underwood, 1997). Before analyses, normality and homoscedasticity of the data were tested using Kolmogorov–Smirnov and Bartlett tests, respectively. All the analyses were performed using the program Statgraphics Plus 5.1.

## Results

### Experimental seawater system

The physical and chemical characteristics of the experimental seawater are summarized in Table 1, and were previously described by Duarte *et al.* (2014) in a parallel study on the calcification rate of mussels. The experimental seawater showed very similar values for salinity at all the temperature/pCO<sub>2</sub> combinations. There was a constant decreasing trend of pH, carbonate contents, and aragonite with the increase in the pCO<sub>2</sub> concentrations at both experimental temperatures.

### Physiological parameters


CR of *M. chilensis* (Figure. 1a) was significantly affected by pCO<sub>2</sub> and temperature (two-way ANOVA; pCO<sub>2</sub>:  $F_{1,12} = 7.72$ ,  $p < 0.01$ ; temperature:  $F_{1,12} = 30.17$ ,  $p < 0.001$ ). Lower values of CR occurred at the highest pCO<sub>2</sub> concentration (1000  $\mu$ atm) compared with the control (380  $\mu$ atm) and with the intermediate concentration of pCO<sub>2</sub> (700  $\mu$ atm). Conversely, CR was significantly higher at 16°C. The interaction between both factors did not have a significant effect ( $p > 0.05$ ) on the CR (temperature \* pCO<sub>2</sub>:  $F_{1,12} = 0.13$ ,  $p > 0.05$ ). A similar pattern was found for the organic ingestion rate, where the highest level of pCO<sub>2</sub> had a negative effect on this physiological process, while a rise in temperature produced an increase in the organic ingestion rate (Figure. 1b).

AE was not affected by temperature; while under the higher concentration of pCO<sub>2</sub> (1000  $\mu$ atm), it was significantly lower than in the control treatment (380  $\mu$ atm CO<sub>2</sub>). Individuals maintained at 700  $\mu$ atm CO<sub>2</sub> exhibited intermediate values, without stating significant differences with the two other treatments (two-way ANOVA; temperature:  $F_{1,12} = 3.00$ ,  $p > 0.05$ ; pCO<sub>2</sub>:  $F_{2,12} = 4.10$ ,  $p < 0.05$ ; temperature \* pCO<sub>2</sub>:  $F_{2,12} = 0.77$ ,  $p > 0.05$ ; Figure. 2a). The absorption rate was affected significantly both by pCO<sub>2</sub> and by temperature, but not for the interaction of both factors. It was significantly lower at the highest pCO<sub>2</sub> concentration compared with the other two treatments; while between 380 and 700  $\mu$ atm pCO<sub>2</sub>, there were no


**Table 1.** Seawater characteristics (mean  $\pm$  SD) used to maintain *M. chilensis* during the experimental period [see Material and methods for details on the temperature and CO<sub>2</sub> values used in the study; from Duarte et al. (2014)].

CO <sub>2</sub> system parameters	Experimental treatments: temperature and CO <sub>2</sub> concentrations							
	12°C		16°C		12°C		16°C	
	380 μatm (current)		700 μatm (2050 year)		1000 μatm (2100 year)			
pH <i>in situ</i> (pH units)	8.057 $\pm$ 0.026	7.996 $\pm$ 0.026	7.835 $\pm$ 0.035	7.776 $\pm$ 0.034	7.673 $\pm$ 0.030	7.618 $\pm$ 0.030		
Salinity (psu)	31.79 $\pm$ 1.88	31.80 $\pm$ 1.66	31.81 $\pm$ 1.90	31.83 $\pm$ 1.72	31.87 $\pm$ 1.87	31.88 $\pm$ 1.66		
TA (μmol kg <sup>-1</sup> )	2156.44 $\pm$ 86.24		2148.63 $\pm$ 90.76		2142.19 $\pm$ 93.23			
pCO <sub>2</sub> <i>in situ</i> (μatm)	371.35 $\pm$ 23.52	439.54 $\pm$ 27.63	656.91 $\pm$ 50.44	772.64 $\pm$ 58.46	977.56 $\pm$ 60.52	1141.90 $\pm$ 69.19		
[CO <sub>3</sub> <sup>2-</sup> ] <i>in situ</i> (μmol kg <sup>-1</sup> )	129.48 $\pm$ 14.65	131.08 $\pm$ 14.74	82.45 $\pm$ 11.24	83.97 $\pm$ 11.32	58.47 $\pm$ 7.92	59.95 $\pm$ 8.00		
Ω <sub>ca</sub>	3.14 $\pm$ 0.33	3.19 $\pm$ 0.33	1.99 $\pm$ 0.25	2.04 $\pm$ 0.25	1.42 $\pm$ 0.18	1.46 $\pm$ 0.18		
Ω <sub>ar</sub>	1.99 $\pm$ 0.22	2.04 $\pm$ 0.22	1.27 $\pm$ 0.17	1.31 $\pm$ 0.17	0.90 $\pm$ 0.12	0.93 $\pm$ 0.12		

TA, total alkalinity; [CO<sub>3</sub><sup>2-</sup>], carbonate ion concentration; Ω<sub>ca</sub>, omega calcite; Ω<sub>ar</sub>, omega aragonite.



**Figure 1.** CR (a) and organic ingestion rate (b) of juvenile individuals of the mussel *M. chilensis* exposed to different combinations of temperature and pCO<sub>2</sub> for 80 d. Values are means  $\pm$  standard error.



**Figure 2.** AE (a) and absorption rate (b) of juvenile individuals of the mussel *M. chilensis* exposed to different combinations of temperature and pCO<sub>2</sub> for 80 d. Values are means  $\pm$  standard error.


significant difference. The absorption rate was significantly higher in mussels exposed to 16°C. (two-way ANOVA; pCO<sub>2</sub>: F<sub>2,12</sub> = 11.20, p < 0.05; temperature: F<sub>1,12</sub> = 28.00, p < 0.01; temperature \* pCO<sub>2</sub>: F<sub>2,12</sub> = 0.28, p > 0.05; Figure. 2b).

The excretion rate of *M. chilensis* was not affected by pCO<sub>2</sub> concentration; however, it was affected significantly by temperature, with higher values at 16°C (two-way ANOVA; pCO<sub>2</sub>: F<sub>2,12</sub> = 0.77, p > 0.05; temperature: F<sub>1,12</sub> = 5.32, p < 0.05; temperature \* pCO<sub>2</sub>: F<sub>2,12</sub> = 0.84, p > 0.05; Figure. 3a).

Oxygen consumption was significantly affected by pCO<sub>2</sub>, but not by temperature or by the temperature \* pCO<sub>2</sub> interaction (two-way

ANOVA; pCO<sub>2</sub>: F<sub>2,12</sub> = 6.10, p < 0.01; temperature: F<sub>1,12</sub> = 0.031, p > 0.05; temperature \* pCO<sub>2</sub>: F<sub>2,12</sub> = 0.26, p > 0.05). Significantly lower values of oxygen uptake were observed in mussels exposed to 1000 μatm pCO<sub>2</sub> compared with those exposed to 380 μatm pCO<sub>2</sub>. Individuals maintained at 700 μatm pCO<sub>2</sub> exhibited intermediate values of oxygen uptake, without stating significant differences with the two other treatments (Figure. 3b).

Two-way ANOVA showed a significant effect of the factors pCO<sub>2</sub> (F<sub>1,12</sub> = 4.38, p < 0.05) and temperature (F<sub>1,12</sub> = 15.58, p < 0.001) on the SFG of *M. chilensis*. However, the interaction between both factors did not have a significant effect (F<sub>1,12</sub> = 0.59, p > 0.05).



**Figure 3.** Ammonia excretion (a), oxygen uptake (b) and SFG (c) of juvenile individuals of the mussel *M. chilensis* exposed to different combinations of temperature and  $p\text{CO}_2$  for 80 days. Values are means  $\pm$  standard error.

SFG was significantly lower at the highest  $p\text{CO}_2$  concentration (1000  $\mu\text{atm}$ ) compared with the control (380  $\mu\text{atm}$ ). Mussels exposed to 700  $\mu\text{atm}$   $p\text{CO}_2$  did not show significantly different values of SFG from the other two  $p\text{CO}_2$  treatments. SFG was significantly higher at 16°C than at 12°C (Figure. 3c).

## Discussion

Our results showed that the rise in temperature from 12 to 16°C in a scenario of elevated  $p\text{CO}_2$  (1000  $\mu\text{atm}$   $p\text{CO}_2$ ) did not increase the sensitivity of the physiological processes measured in *M. chilensis*. On the contrary, it was observed that the increase in temperature induced a positive effect on the physiological processes associated with increased energy. In turn, the increase of  $p\text{CO}_2$  adversely affected this species by reducing the energy intake, which resulted

in significantly lower values of SFG at 1000  $\mu\text{atm}$   $p\text{CO}_2$  at both temperatures, compared with 380  $\mu\text{atm}$   $p\text{CO}_2$ .

Contrasting results have been described in previous studies on the combined effects of warming and elevated  $p\text{CO}_2$  levels on different groups of marine invertebrates. Catarino *et al.* (2012) demonstrated that ocean warming and acidification antagonistically interacted on oxygen uptake of the sea urchin *Paracentrotus lividus*. Byrne and Przeslawski (2013) reviewed the combined effect of ocean warming and high  $p\text{CO}_2$  levels in >20 species of invertebrates. Their results showed that antagonistic effects were common, where ocean warming reduces the negative effects of  $p\text{CO}_2$ , unlike synergies, which were less common. Previous studies have shown that feeding activity in molluscs can increase within the thermal window of each species and drastically reduced when the temperature exceeds the limit of thermal tolerance (Navarro *et al.*, 2002; Peck *et al.*, 2009). Wang *et al.* (2015) found that increasing temperature significantly reduced CR of the mussel *Mytilus coruscus*, but elevated levels of  $p\text{CO}_2$  did not show an additive or antagonistic effect with temperature. In our study, temperature positively affected CR, increasing from 12 to 16°C at all experimental  $p\text{CO}_2$  levels. This can be explained by the experimental temperature range used, which is within the thermal window (9–16°C) of the studied population of *M. chilensis* (Navarro and Jaramillo, 1994). The negative effect of elevated concentrations of  $p\text{CO}_2$  on the feeding rate of *M. chilensis* observed in our study has been also described for other species of bivalves. Fernandez-Reiriz *et al.* (2011) observed a reduction in the feeding rate in the clam *Ruditapes decussatus* at the highest experimental  $p\text{CO}_2$  concentration (pH 7.48). Navarro *et al.* (2013) also described negative effects of elevated levels of  $p\text{CO}_2$  (pH 7.57) on the feeding rate of the mussel *M. chilensis*.

There are few studies on the combined effects of ocean warming and high  $p\text{CO}_2$  levels on AE of marine organisms. Zhang *et al.* (2015) found that AE of the gastropod *Nassarius conoidalis* was not significantly affected at the beginning of the experimental period (day 2) by temperature,  $p\text{CO}_2$  level, nor the interaction between temperature and  $p\text{CO}_2$  levels. However, on day 30, the temperature negatively affected AE when this gastropod was exposed at the highest  $p\text{CO}_2$  concentration. Wang *et al.* (2015) described low values for AE (20–45%) on the mussel *M. coruscus* with no significant differences between three different pH levels (8.1, 7.7, and 7.3). Our results showed that  $p\text{CO}_2$  levels significantly affected the AE of *M. chilensis*, but this effect was not registered for neither temperature nor the interaction between temperature and  $p\text{CO}_2$ . In a previous study, Navarro *et al.* (2013) reported that elevated  $p\text{CO}_2$  levels (pH 7.57) reduced significantly the AE of *M. chilensis*, indicating possible deficiencies in the functioning of the digestive system under conditions of seawater acidification.

Similar to other studies on marine bivalves (Bayne and Newell, 1983; Velasco and Navarro, 2005), ammonia excretion of *M. chilensis* represented a low amount of the energy absorbed, with a minimal impact on the energy balance. Ammonia excretion was only affected by temperature, and the increase observed in mussels exposed to the highest temperature (16°C) could be explained by the larger amount of food ingested and absorbed under these conditions. Navarro *et al.* (2013) found similar values for ammonia excretion in juveniles of *M. chilensis* exposed to a wide range of  $p\text{CO}_2$  levels (7.91–7.57).

There is evidence of the capability of marine organisms to acclimate the metabolic rate within certain range of temperature (thermal window), being a species-specific response (Pörtner *et al.*, 2005; Pörtner, 2008, 2010; Ezgeta-Balic *et al.*, 2011). However, at high temperatures, beyond the range of thermal tolerance of a species, a

drastic decline occurs in oxygen consumption and an increase in the anaerobic metabolic pathways (Jansen *et al.*, 2007). In our study, oxygen consumption was not affected by temperature. This can be explained because the mussels were exposed to a temperature range that the animals experience in their natural environment. However, temperatures over the thermal window naturally could have a negative impact on oxygen consumption of *M. chilensis*. Several studies have described metabolic depression in different species of marine organisms at elevated  $p\text{CO}_2$  concentrations, caused by the low capacity to compensate for disturbances in extracellular ion and acid–base status (Michaelidis *et al.*, 2005; Pörtner, 2008). Our study showed a depletion of oxygen consumption when mussels were exposed to the highest  $p\text{CO}_2$  level, suggesting that *M. chilensis* also is not able to cope with high  $p\text{CO}_2$  levels in the seawater.

Different physiological processes related with gain and expenditure of energy largely depend on an organism's environmental temperature (Bayne and Newell, 1983). Populations that can maintain their aerobic capacity at warmer temperatures have a higher thermal tolerance, and are thereby predicted to persist longer as the temperature increases (Pörtner, 2001; Gardiner *et al.*, 2010). SFG represents the integration of all these processes varying as a function of temperature (Ezgeta-Balic *et al.*, 2011). Wang *et al.* (2015), studying the combined effects of seawater temperature and pH on the mussel *M. coruscus*, described positive values of SFG for all treatments, with no significant effects of pH. However, temperature showed a significant effect on the SFG, being reduced from 25 to 30°C. Ezgeta-Balic *et al.* (2011) recorded negative values of SFG for the mussel *Modiolus barbatus* when temperature rises from 26 to 28°C. Anestis *et al.* (2010), when studying the response of the mussel *Mytilus galloprovincialis* to increasing seawater temperature, found that the SFG values became negative at temperatures higher than 24°C, associated with a reduction in CR. Contrary to these authors, the present study showed a significant positive effect of temperature on SFG at all  $p\text{CO}_2$  levels. This might be explained because the experimental mussels were exposed to temperatures experienced in their natural environment, which are within the range of thermal tolerance of the species although the highest temperature (16°C) is recorded in a narrow temporal window during summer (Navarro and Jaramillo, 1994). The increase of the SFG observed in the present study matches well with the significant increase of CR and absorption rate from 12 to 16°C. Oxygen uptake was not affected by increasing temperature, suggesting that metabolic activity is maintained constant within the thermal window of *M. chilensis*. Similar conclusions were reported by Findlay *et al.* (2010) to explain the lack of differences between the growth rates of *Semibalanus balanoides* exposed to a range of temperatures experienced in their natural environment. The zone of the Reloncaví Sound, located 50 km east from the place where the experimental mussels were collected (Bay of Huelmo), has been described as an environment with a low calcium carbonate saturation and even “corrosive” when occurring in waters with low salinity and alkalinity (Alarcón *et al.*, 2015). This can explain the lack of significant differences in some of the physiological responses between the 380 and 700  $p\text{CO}_2$  levels, suggesting that mussels from this area might be pre-adapted to natural pH variability.

There are still few studies on the combined effect of  $p\text{CO}_2$  with other environmental stressors. Gazeau *et al.* (2014) found that somatic and shell growth of *M. galloprovincialis* did not appear very sensitive to OA and warming during most of the experiment. However, growth was significantly reduced after summer in the lowered pH treatment. According to these authors, there is a

progressive insufficiency in acid–base regulation capacity, which was also consistent with shell net dissolution observed in the mussels (Gazeau *et al.*, 2014). Other studies described that growth might be significantly reduced under medium- and long-term exposure to elevated  $p\text{CO}_2$  levels (Beniash *et al.*, 2010; Navarro *et al.*, 2013; Duarte *et al.*, 2014). Both shell and soft tissue growth of the oyster *C. virginica* were reduced when exposed to high  $p\text{CO}_2$  levels (pH 7.5; Beniash *et al.*, 2010). Similarly, Duarte *et al.* (2014) found that net rate of calcium deposition and total weight of the mussel *M. chilensis* were not significantly affected by temperature, but were negatively affected by high  $p\text{CO}_2$  levels. Our results suggest that a rise in temperature within the natural range experienced by juveniles *M. chilensis* generates a positive effect on the processes related with energy gain from the environment (i.e. feeding and absorption) to be allocated for growth. In turn, the increase in 1000  $\mu\text{atm } p\text{CO}_2$ , independent of temperature, adversely affects this species, with significantly reduced energy allocated to growth (SFG) compared with the control treatment (380  $\mu\text{atm } p\text{CO}_2$ ). Our results are interpreted as answers to short-term toxicity challenges with temperature and  $\text{CO}_2$  as environmental factors. On the basis of these, inferences about environmental changes that would slowly occur over the next 100–200 years must be made with appropriate caution. This is because animals used for testing would vary in geno- and phenotypes.

The effect of elevated  $p\text{CO}_2$  on marine organisms can be negative (Gazeau *et al.*, 2014), reduced, or absent in the presence of high temperature (Pörtner and Farrel, 2008; Duarte *et al.*, 2014). This is because the effect of warming will depend on the experimental temperature range and thermal window of the species under study. Thus, the studies related to the combined effect of ocean warming and elevated concentrations of  $p\text{CO}_2$  must consider the thermal windows of each species to avoid contradictory interpretations. Future experiments with different levels of temperatures should be designed in a way that will allow greater predictive power, selecting temperature treatments that will be physiologically realistic to the animal's thermal windows and to future climate scenarios.

## Acknowledgements

The authors are grateful to Elizabeth Encalada and Alejandro Ortiz for their assistance during the laboratory and fieldwork. Special thanks are due to the reviewers for very constructive comments on the manuscript. This study was mainly funded by the Programa de Investigación Asociativa, PIA CONICYT-CHILE (Grant Anillos ACT-132). Additional support came from FONDECYT Grant 1120470 to JMN and from Center FONDAP-CONICYT IDEAL 15150003. CAV is supported by Red Doctoral (REDOC.CTA) MINEDUC project UCO1202 at Universidad de Concepción.

## References

- Alarcón, E., Valdés, N., and Torres, R. 2015. Saturación del carbonato de calcio en un área de cultivo de mitílidos en el Seno Reloncaví, Patagonia norte, Chile. *Revista Chilena de Historia Natural*, 43: 277–281.
- Anestis, A., Pörtner, H. O., Karagiannis, D., Angelidis, P., Staikou, A., and Michaelidis, B. 2010. Response of *Mytilus galloprovincialis* (L.) to increasing seawater temperature and to marfanosis: Metabolic and physiological parameters. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology*, 156(1): 57–66.
- Bayne, B. L., and Newell, R. C. 1983. Physiological energetics of marine molluscs. In *The Mollusca* Vol. 4. *Physiology*. Part 1, pp. 407–515. Ed. by K. M. Wilbur. Academic Press, New York.

- Beniash, E., Ivanina, A., Lieb, N. S., Kurochkin, I., and Sokolova, I. M. 2010. Elevated level of carbon dioxide affects metabolism and shell formation in oysters *Crassostrea virginica*. *Marine Ecology Progress Series*, 419: 95–108.
- Byrne, M. 2011. Impact of ocean warming and ocean acidification on marine invertebrate life history stages: Vulnerabilities and potential for persistence in a changing ocean. *Oceanography and Marine Biology: An Annual Review*, 49: 1–42.
- Byrne, M., and Przeslawski, R. 2013. Multistressor impacts of warming and acidification of the ocean on marine invertebrates' life histories. *Integrative and Comparative Biology*, 53: 582–596.
- Caldeira, K., and Wickett, M. E. 2003. Oceanography: Anthropogenic carbon and ocean pH. *Nature*, 425: 365.
- Catarino, A. I., Bauwens, M., and Dubois, P. 2012. Acid-base balance and metabolic response of the sea urchin *Paracentrotus lividus* to different seawater pH and temperatures. *Environmental Science and Pollution Research International*, 19: 2344–2353.
- Conover, R. J. 1966. Assimilation of organic matter by zooplankton. *Limnology and Oceanography*, 11: 338–345.
- Coughlan, J. 1969. The estimation of filtering rate from the clearance of suspensions. *Marine Biology*, 2: 356–358.
- Dickson, A. G., and Millero, F. J. 1987. A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep-Sea Research Part A: Oceanographic Research Papers*, 34: 1733–1743.
- Duarte, C., Jaramillo, E., Contreras, H., and Figueroa, L. 2006. Community structure of the macroinfauna in the sediments below and intertidal mussel bed (*Mytilus chilensis*) (Hupe) of southern Chile. *Revista Chilena de Historia Natural*, 79: 353–368.
- Duarte, C., Navarro, J. M., Acuña, K., Torres, R., Manriquez, P. H., Lardies, M. A., Vargas, C. A., et al. 2014. Combined effects of temperature and ocean acidification on the juvenile individuals of the mussel *Mytilus chilensis*. *Journal of Sea Research*, 85: 308–314.
- Elliot, J. M., and Davison, W. 1975. Energy equivalents of oxygen consumption in animal energetics. *Oecologia*, 19: 195–201.
- Ezgeta-Balic, D., Rinaldi, A., Peharda, M., Prusina, I., Montalto, V., Niceta, N., and Sarà, G. 2011. An energy budget for the subtidal bivalve *Modiolus barbatus* (Mollusca) at different temperatures. *Marine Environmental Research*, 71: 79–85.
- Fabry, V. J., Seibel, B. A., Feely, R. A., and Orr, J. C. 2008. Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES Journal of Marine Science*, 65: 414–432.
- Feely, R. A., Sabine, C. L., Lee, K., Berelson, W., Kleypas, J., Fabry, V. J., and Millero, F. J. 2004. Impact of anthropogenic CO<sub>2</sub> on the CaCO<sub>3</sub> system in the oceans. *Science*, 305: 362–366.
- Fernandez-Reiriz, M. J., Range, P., Álvarez-Salgado, X. A., and Labarta, U. 2011. Physiological energetics of juvenile clams *Ruditapes decussatus* in a high CO<sub>2</sub> coastal ocean. *Marine Ecology Progress Series*, 433: 97–105.
- Findlay, H. S., Kendall, M. A., Spicer, J. I., and Widdicombe, S. 2010. Post-larval development of two intertidal barnacles at elevated CO<sub>2</sub> and temperature. *Marine Biology*, 157: 725–735.
- Gardiner, N. M., Munday, P. L., and Nilsson, G. E. 2010. Counter-gradient variation in respiratory performance of coral reef fishes at elevated temperatures. *PLoS ONE*, 5: e13299.
- Gazeau, F., Allouane, S., Bock, C., Bramanti, L., Lopez Correa, M., Gentile, M., Hirse, T., et al. 2014. Impact of ocean acidification and warming on the Mediterranean mussel (*Mytilus galloprovincialis*). *Frontiers Marine Science*, 1: 62.
- Gobler, C. J., and Talmage, S. C. 2014. Physiological response and resilience of early life-stage Eastern oysters (*Crassostrea virginica*) to past, present and future ocean acidification. *Conservation Physiology*, 2. doi:10.1093/conphys/cou004.
- Godbold, J. A., and Solan, M. 2013. Long-term effects of warming and ocean acidification are modified by seasonal variation in species responses and environmental conditions. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences*, 368: 20130186.
- Gooding, R. A., Harley, C. D. G., and Tang, E. 2009. Elevated water temperature and carbon dioxide concentration increase the growth of a keystone echinoderm. *Proceedings of the National Academy of Sciences*, 106: 9316–9321.
- Haraldsson, C., Anderson, L. G., Hassellöv, M., Hulth, S., and Olsson, K. 1997. Rapid, high-precision potentiometric titration of alkalinity in ocean and sediment pore waters. *Deep-Sea Research Part 1: Oceanographic Research Papers*, 44: 2031–2044.
- Harley, C. D., Randall Hughes, A., Hultgren, K. M., Miner, B. G., Sorte, C. J., Thornber, C. S., Rodriguez, L. F., et al. 2006. The impacts of climate change in coastal marine systems. *Ecology Letters*, 9: 228–241.
- Hiebenthal, C., Philipp, E. E. R., Eisenhauer, A., and Wahl, M. 2013. Effects of seawater pCO<sub>2</sub> and temperature on shell growth, shell stability, condition and cellular stress of Western Baltic Sea *Mytilus edulis* (L.) and *Arctica islandica* (L.). *Marine Biology*, 160: 2073–2087.
- IPCC. 2013. The Physical Science Basis. Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge.
- Ivanina, A. V., Dickinson, G. H., Matoo, O. B., Bagwe, R., Dickinson, A., Beniash, E., and Sokolova, I. M. 2013. Interactive effect of elevated temperature and CO<sub>2</sub> levels on energy metabolism and biomineralization of marine bivalves *Crassostrea virginica* and *Mercenaria mercenaria*. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology*, 166: 101–111.
- Jansen, J. M., Pronker, A. E., Kube, S., Sokolowski, A., Sola, J. C., Marquiegui, M. A., Shiedek, D., et al. 2007. Geographic and seasonal patterns and limits on the adaptive response to temperature of European *Mytilus* spp. and *Macoma balthica* populations. *Oecologia*, 154: 23–34.
- Lewis, E., and Wallace, D. 1998. Program Developed for CO<sub>2</sub> System Calculations. Carbon Dioxide Information Analysis Center, Oak Ridge, TN, USA.
- Mehrbach, C., Culbertson, C. H., Hawley, J. E., and Pytkowicz, R. M. 1973. Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnology and Oceanography*, 18: 897–907.
- Meier, H. E. M. 2006. Baltic Sea climate in the late twenty-first century: A dynamical downscaling approach using two global models and two emission scenarios. *Climate Dynamics*, 27: 39–68.
- Metzger, R., Sartoris, F. J., Langenbuch, M., and Pörtner, H. O. 2007. Influence of elevated CO<sub>2</sub> concentrations on thermal tolerance of the edible crab *Cancer pagurus*. *Journal of Thermal Biology*, 32: 144–151.
- Michaelidis, B., Ouzounis, C., Palaras, A., and Pörtner, H. O. 2005. Effects of long-term moderate hypercapnia on acid-base balance and growth rate in marine mussels *Mytilus galloprovincialis*. *Marine Ecology Progress Series*, 293: 109–118.
- Navarro, J. M., and Jaramillo, R. 1994. Evaluación de la oferta alimentaria natural disponible a organismos filtradores en la Bahía de Yaldad (43–08; 73–44'), sur de Chile. *Revista de Biología Marina*, 29: 57–75.
- Navarro, J. M., Leiva, G. E., Gallardo, C. S., and Varela, C. 2002. Influence of diet and temperature on physiological energetics of *Chorus giganteus* (Gastropoda:Muricidae) during reproductive conditioning. *New Zealand Journal of Marine and Freshwater Research*, 36: 321–332.
- Navarro, J. M., Leiva, G. E., Martínez, G., and Aguilera, C. 2000. Interactive effects of diet and temperature on the scope for growth of the scallops *Argopecten purpuratus* during reproductive conditioning. *Journal of Experimental Marine Biology and Ecology*, 247: 67–83.
- Navarro, J. M., Torres, R., Acuña, K., Duarte, C., Manriquez, P. H., Lardies, M., Lagos, N. A., et al. 2013. Impact of medium-term

- exposure to elevated  $p\text{CO}_2$  levels on the physiological energetic of the mussel *Mytilus chilensis*. *Chemosphere*, 90: 1242–1248.
- Ohle, W. 1953. Die chemische und elektrochemische Bestimmung des molekular gelösten Sauerstoffs der Binnengewässer. *Mitteilungen Internationale Vereinigung der Limnologie*, 3: 1–44.
- Peck, L. S., Massey, A., Thorne, M. A., and Clark, M. S. 2009. Lack of acclimation in *Ophionotus victoriae*: Brittle stars are not fish. *Polar Biology*, 32: 399–402.
- Philippart, C. J. M., Van Aken, H. M., Beukema, J. J., Bos, O. G., Cadée, G. C., and Dekker, R. 2003. Climate-related changes in recruitment of the bivalve *Macoma balthica*. *Limnology and Oceanography*, 48: 2171–2185.
- Pörtner, H. O. 2001. Climate change and temperature-dependent biogeography: Oxygen limitation of thermal tolerance in animals. *Naturwissenschaften*, 88: 137–146.
- Pörtner, H. O. 2008. Ecosystem effects of ocean acidification in times of ocean warming: A physiologist's view. *Marine Ecology Progress Series*, 373: 203–217.
- Pörtner, H. O. 2010. Oxygen- and capacity-limitation of thermal tolerance: A matrix for integrating climate-related stressor effects in marine ecosystems. *The Journal of Experimental Biology*, 213: 881–893.
- Pörtner, H. O., and Farrel, A. P. 2008. Physiology and climate change. *Science*, 322: 690–692.
- Pörtner, H. O., Langenbuch, M., and Michaelidis, B. 2005. Synergistic effects of temperature extremes, hypoxia, and increases in  $\text{CO}_2$  on marine animals: From Earth history to global change. *Journal of Geophysical Research*, 110: 1–15.
- Pörtner, H. O., Langenbuch, M., and Reipschläger, A. 2004. Biological impact of elevated ocean  $\text{CO}_2$  concentrations: Lessons from animal physiology and Earth history. *Journal of Oceanography*, 60: 705–718.
- Schiedek, D., Sundelin, B., Readman, J. W., and Macdonald, R. W. 2007. Interactions between climate change and contaminants. *Marine Pollution Bulletin*, 54: 1845–1856.
- Solórzano, L. 1969. Determination of ammonia in natural waters by the phenylhypochlorite method. *Limnology and Oceanography*, 14: 799–801.
- Somero, G. N. 2002. Thermal physiology and vertical zonation of intertidal animals: Optima, limits and costs of living. *Integrative and Comparative Biology*, 42: 780–789.
- Torres, R., Manriquez, P. H., Duarte, C., Navarro, J. M., Lagos, N. A., Vargas, C. A., and Lardies, M. A. 2013. Evaluation of a semi-automatic system for long-term seawater carbonate chemistry manipulation. *Revista Chilena de Historia Natural*, 86: 443–451.
- Underwood, A. J. 1997. *Experiments in Ecology: Their Logical Design and Interpretation Using Analysis of Variance*. Cambridge University Press, Cambridge.
- Vargas, C. A., De la Hoz, M., Aguilera, V., San Martín, V., Manriquez, P. H., Navarro, J. M., Torres, R., *et al.* 2013.  $\text{CO}_2$ -driven ocean acidification reduces larval feeding efficiency and change food selectivity in the mollusk *Concholepas concholepas*. *Journal of Plankton Research*, 35: 1059–1068.
- Velasco, L. A., and Navarro, J. M. 2005. Feeding physiology of two bivalves under laboratory and field conditions in response to variable food concentrations. *Marine Ecology Progress Series*, 291: 115–124.
- Walther, G. R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T., Fromentin, J.-M., *et al.* 2002. Ecological responses to recent climate change. *Nature*, 416: 389–395.
- Wang, Y., Li, L., Hu, M., and Lu, W. 2015. Physiological energetic of the thick shell mussel *Mytilus coruscus* exposed to seawater acidification and thermal stress. *Science of the Total Environment*, 1: 261–272.
- Whyte, J. 1987. Biochemical composition and energy content of six species of phytoplankton used in mariculture of bivalves. *Aquaculture*, 60: 231–241.
- Widdicombe, S., and Spicer, J. I. 2008. Predicting the impact of ocean acidification on benthic biodiversity: What can animal physiology tell us? *Journal of Experimental Marine Biology and Ecology*, 366: 187–197.
- Zhang, H., Shin, P. K. S., and Cheung, S. G. 2015. Physiological responses and scope for growth upon medium-term exposure to the combined effects of ocean acidification and temperature in a subtidal scavenger *Nassarius conoidalis*. *Marine Environmental Research*, 106: 51–60.

Handling editor: Howard Browman